

Symplectic Generating Functions and Moyal Products

PAVOL ŠEVERA^{1,2}

¹*Section de Mathématiques, Université de Genève, Rue du Lièvre 2-4, 1211 Genève, Switzerland. e-mail: paval.severa@gmail.com*

²*Department of Theoretical Physics, Comenius University, Mlynská dolina F2, 84248 Bratislava, Slovakia*

Received: 28 September 2005


Published online: 22 February 2006

Abstract. We introduce an affine-invariant version of generating functions of symplectic transformations of affine symplectic spaces, together with a generalization for other symmetric symplectic spaces. The composition of these functions has a nice connection with the Moyal product.

Mathematics Subject Classifications (2000). 53D22, 53C35.

Key words. symplectomorphisms, generating functions, Moyal product, symmetric spaces.


Let us start with the simplest, but probably the most useful result of this note. Suppose A is an affine symplectic space. There is an affine-invariant way in which functions on A generate symplectic transformations of A . Namely, let H be a function on A . At any point $x \in A$ we take the vector u_x defined by $d_x H = \omega(\cdot, u_x)$ (where ω is the symplectic form) and substitute it in A so that x lies in its middle. Then the map Φ_H sending the tails of u_x 's to their heads is a symplectic transformation:



Notice that for infinitesimal Hamiltonians, this is the usual infinitesimal transformation generated by the Hamiltonian H . The map $H \mapsto \Phi_H$ is a kind of Cayley transform: choosing an origin in A to make it to a vector space, and restricting ourselves to H 's which are quadratic forms, we get the usual Cayley transform $\mathfrak{sp} \rightarrow Sp$.

Symplectic transformation can, of course, be composed. The corresponding composition of generating functions is

$$H(x) = H_1(x_1) + H_2(x_2) + \text{symplectic area of } \triangle PQR:$$




This composition law is closely related with the integral kernel of the Moyal product,

$$K(x_1, x_2, x) = \exp(\sqrt{-1} \times \text{symplectic area of } \triangle PQR/\hbar):$$

the function $\exp(\sqrt{-1}H/\hbar)$ is the leading oscillatory part of the Moyal product of $\exp(\sqrt{-1}H_1/\hbar)$ with $\exp(\sqrt{-1}H_2/\hbar)$.

The composition law appeared previously in [2], which was inspired earlier in [1], which should refer to even earlier [5]. More on the connection with quantization can be found in [4].

Let us prove our claims. The graph of any symplectic transformation of A is a Lagrangian submanifold of $\bar{A} \times A$. For each point $x \in A$ the symmetry with respect to x is a symplectic transformation. Their graphs, together with the graph of the identity, can be used to identify $\bar{A} \times A$ with T^*A :




Explicitly (as one immediately sees from the picture), a pair $(P, Q) \in \bar{A} \times A$ corresponds to $((P + Q)/2, \omega(\cdot, Q - P)) \in T^*A$. Here the vector-and-its-midpoint picture emerges.

The correspondence between generating functions and symplectic transformations is clear now: dH is a Lagrangian submanifold of T^*A , and therefore of $\bar{A} \times A$; if it is the graph of some transformation of A , the transformation is symplectic.

Let us explain where the composition law for generating functions come from. The space $\Gamma = \bar{A} \times A$ is a symplectic groupoid (the pair groupoid of A). The graph of its multiplication is a Lagrangian submanifold in $\Gamma \times \Gamma \times \bar{\Gamma}$; using the identification of T^*A with $\Gamma = \bar{A} \times A$, it should be given by a closed 1-form on $A \times A \times A$. This 1-form is the differential of the function $(x_1, x_2, x) \mapsto$ symplectic area of $\triangle PQR$ (this was observed in much greater generality by Alan Weinstein in [6]; we shall need his more general statement later). The composition law and its connection with Moyal product are now clear.

For the fun of it, let us make a similar construction, replacing A by the sphere S^2 with its area symplectic form. Again symmetry with respect to a point of S^2 (i.e., rotation of the sphere by π around the axis passing through that point) is a symplectic transformation of the sphere, therefore we locally have a similar identification between $\bar{S}^2 \times S^2$ and T^*S^2 ; more precisely, there is an isomorphism between the subset of covectors in T^*S^2 of length less than 2 and $\bar{S}^2 \times S^2$ with erased pairs of antipodal points. Explicitly, to a non-antipodal pair (P, Q) we associate a $u \in TS^2$ (and thus, via ω , a point in T^*S^2) as on the picture:



Here x is the midpoint of the short geodesic arc PQ and $u \in T_x S^2$ appears by the orthogonal projection of the arc. This picture can be derived from the famous theorem of Archimedes, claiming that a certain map between the cylinder and sphere is area-preserving.

As a result, we have a similar picture of generating functions: for a function H on S^2 and any point $x \in S^2$ we take the vector u_x defined by $d_x H = \omega(\cdot, u_x)$, place it into the tangent plane $T_x S^2$ so that x is in its middle and project it into the sphere; Φ_H maps P to Q . Composition rule looks as before, only the triangles are spherical now (as follows again from [6]).

Generally, this picture works with no changes for an arbitrary symmetric symplectic space M : we embed M into an affine space A (just as S^2 is embedded into \mathbb{R}^3) and extend the symmetry of M with respect to any point $x \in M$ to an involution σ_x of A . Then we project M to $T_x M$ in the direction of A^{σ_x} (just as we used the orthogonal projection of the sphere to its tangent planes). The generation of symplectic transformations of M by functions on M , and the composition law for generating functions, are then the same as for the sphere (this was also noticed in [3], in a somewhat different way).

Here are the details: Since M is a homogeneous space of some Lie group G , we can find a moment map $M \rightarrow \mathfrak{g}^*$ (the map is possibly non-equivariant and defined only on a covering of M ; in fact, being a homogeneous symplectic space, M is, up to covering, a coadjoint orbit of a central extension of G). This is the immersion $M \rightarrow A$ we need. Now, since M is a symmetric space, it is (up to covering) G/G^σ , where σ is an involutory automorphism of G . Let $\mathfrak{g} = \mathfrak{g}^\sigma \oplus \mathfrak{p}$ be the decomposition of \mathfrak{g} to ± 1 eigenspaces of $d\sigma$; to make G/G^σ to a symmetric symplectic space, we have to choose a \mathfrak{g}^σ -invariant symplectic form on \mathfrak{p} . If $x \in M$ is fixed by G^σ (for other x values we have to conjugate σ with the corresponding inner automorphism of G), $T_x M$ is \mathfrak{p}^* translated to x ; we project M to $T_x M$ in the direction of $(\mathfrak{g}^*)^\sigma$.

To finish, we have to prove that these projections $M \rightarrow T_x M$ actually give us a local symplectomorphism between $\bar{M} \times M$ and T^*M . Let us denote the projection by $\pi : M \rightarrow T_x M$. Let us choose $t \in \mathfrak{p}$ and $u \in T_x M = \mathfrak{p}^*$; t gives us a vector field on M , its value at $y \in M$ will be denoted by t_y . We have to prove that for any $\lambda \in \mathbb{R}$ we have

$$\omega_x(u, t_x) = \omega_{\pi^{-1}(\lambda u)}(\pi_*^{-1}u, t_{\pi^{-1}(\lambda u)});$$

this will ensure that the fundamental 1-form on T^*M is mapped to the primitive of the symplectic form on $\bar{M} \times M$ given by its Lagrangian fibration. To prove the formula, we just notice that its RHS is equal to $\langle \pi_*^{-1}u, t \rangle$, which is in turn equal to $\langle u, t \rangle$ (since we project in a direction that is in the kernel of t), which is the LHS.

Acknowledgements

I would like to thank Alan Weinstein for drawing my attention to his work [6] after I posted a first version of this note to the arXiv (math.SG/9905052) in May 1999. I would also like to thank to the referee for suggesting all the other references (though it basically means that there is nothing new in this letter).

References

1. Marinov, M.S.: An alternative to the Hamilton–Jacobi approach in classical mechanics. *J. Phys A, Math. Gen.* **12**, 31–47 (1979)

2. Ozorio de Almeida, A.: On the symplectically invariant variational principle and generating functions. *Proc R Soc London A* **431**, 403–417 (1990)
3. de M Rios, P., Ozorio de Almeida, A.: A variational principle for actions on symmetric symplectic spaces. *J Geom Phys* **51**, 404–441 (2004)
4. de M Rios, P., Tuynman, G.M.: On Weyl quantization from geometric quantization. [math-ph/0201044](https://arxiv.org/abs/math-ph/0201044)
5. Weinstein, A.: On the invariance of Poincare's generating function for canonical transformations. *Invent. Math.* **16**, 202–213 (1972)
6. Weinstein, A.: Traces and triangles in symmetric symplectic spaces. *Symplectic geometry and quantization* (Sanda and Yokohama 1993), pp. 261–270, *Contemporary Mathematics* 179, American Mathematical Society, Providence (1994)